

Aharonov Anandan Phase for the Quasi Exactly Solvable Bose Systems

ANIRBAN PATHAK

*Department of Physics
Visva Bharati
Santiniketan-731235
India*

Abstract

An extended notion of quasi-exactly solvable potential model is used here to treat quasi exactly solvable (QES) Bose systems. We report an analytic expression for the Aharonov Anandan non-adiabatic geometric phase for the QES Bose system in general. The generalized expression is then used to study some particular cases of physical interest and we observe that the geometric phase can be tuned.

1 Introduction:

Phase is responsible for all the interference phenomenon we observe in classical and quantum physics. So people have studied the properties of the quantum phase from different contexts since the beginning of quantum mechanics[1]. But the interest increased considerably in the recent past when Berry published his famous paper [2]. He showed that the phase acquired by the quantum system during a cyclic evolution under the action of an adiabatically varying Hamiltonian is a sum of two parts. The first part is dynamical in nature and the second is geometric. Later on Aharonov and Anandan generalized the idea of the Berry phase and they defined a geometric phase factor for any cyclic evolution of quantum system [3]. The existence of geometric phase is found in many areas of physics. The examples are from the quantum Hall effect to the Jahn-Teller effect and from the spin orbit interaction to quantum computation [1,4,5].

Physicists have always tried to understand Nature in terms of simple models and the simple harmonic oscillator (SHO) is probably the most important one among those models. The SHO model can be used to understand a wide variety of physical phenomena ranging from problems in Newtonian mechanics to those in quantum field theory. However, for a real physical problem one has to incorporate anharmonicity in the model Hamiltonian. For example, it is related to quantum Bose models with interaction or self-interaction. The Schrodinger equation for these anharmonic Bose oscillators are not exactly solvable. However, Dolya and Zaslavaskii [6] have recently shown that the notion of the quasi exactly solvable (QES) models can be extended for anharmonic Bose oscillators.

They have reported that a class of Hamiltonian, often faced by us in molecular physics, theory of magnetism and other branches of physics is QES. This fact provoked us to study the possibilities of observing non-adiabatic geometric phase or Aharonov-Anandan phase for the quasi exactly solvable Bose systems. In the present paper we have found an analytic expression in closed form for the Aharonov-Anandan phase for the QES Bose systems characterized by the Hamiltonian defined by Dolya and Zaslavaskii [6]. The generalized expression is then used to study a particular case of physical interest.

In section 2 we give a short introduction to the Aharonov-Anandan phase[3,7]. We obtain a generalized expression for geometric phase of QES Bose systems in section 3 and use that expression to study some particular cases of physical interest in the subsequent subsection. We devote section 4 for discussions and conclusions.

2 Aharonov-Anandan Phase

In a quantum mechanical description of any system, one has a complex Hilbert space $\mathcal{H}$ whose nonzero vectors represent states of the system. A vector $|\psi\rangle$ in $\mathcal{H}$ and any complex multiple of it (i.e $c|\psi\rangle$) represent the same state in the Hilbert space $\mathcal{H}$. This arbitrariness in the representation of states can be reduced by imposing the normalization condition

$$\langle \psi | \psi \rangle = 1. \quad (1)$$

There still remains an arbitrariness of a phase factor because a normalized vector $|\psi\rangle$ and the vectors $\alpha|\psi\rangle$ represent the same state in Hilbert space provided that the modulus of α is unity (i.e $|\alpha| = 1$). The collection of all vectors $\alpha|\psi\rangle$ with $|\psi\rangle$, a fixed normalized vector and α taking all possible complex values of modulus one is called a unit ray and is denoted by $|\psi\rangle$ [8]. More general rays consist of a collection of all vectors of the form $c|\psi\rangle$. So rays represent vectors without any arbitrariness of phase.

To understand the geometric phase let us start from a simple situation in which a state vector $|\psi\rangle$ evolves cyclically in the Hilbert space $\mathcal{H}$ and after a complete period T returns to the same state vector with a different phase which may be written as $\alpha|\psi\rangle = \exp(i\phi)|\psi\rangle$. Therefore,

$$|\psi(T)\rangle = \exp(i\phi)|\psi(0)\rangle, \quad (2)$$

where ϕ is the total phase which has two parts. The first part is the dynamical phase and the second is the geometric phase. We are interested in the later type of phase which can be found out by subtracting the dynamical phase part from the total phase.

The cyclic evolution of the state vector may be described by a curve C in $\mathcal{H}$ because $|\psi\rangle$ and $\alpha|\psi\rangle$ are two different points in $\mathcal{H}$. Since there are infinitely many possible values of α so there are infinitely many curves C in $\mathcal{H}$ which describes the same cyclic evolution. In the projective Hilbert space of rays P

or in the ray-representative space all these points are represented by the same ray (i.e by the same point). So we will have a single closed curve $\tilde{C}$ in the ray representation space P corresponding to the infinitely many possible curves in the usual Hilbert space. As there is no arbitrariness of phase in P so in the interval $[0, T]$ we must have

$$|\tilde{\psi}(T)\rangle = |\tilde{\psi}(0)\rangle. \quad (3)$$

Aharonov and Anandan[3] exploited this property of single valuedness of the unit ray in the ray representative space P to establish the presence of a geometric phase for all cyclic evolutions. They started by supposing that the normalized state $|\psi(t)\rangle \in \mathcal{H}$ evolves according to the Schrödinger equation

$$H(t)|\psi(t)\rangle = i\hbar \left(\frac{d}{dt} |\psi(t)\rangle \right). \quad (4)$$

We can always define a state vector $|\tilde{\psi}(t)\rangle$ in P which is equivalent to $|\psi(t)\rangle$ as

$$|\tilde{\psi}(t)\rangle = \exp(-if(t))|\psi(t)\rangle. \quad (5)$$

Now substituting (2 and 5) in (3) we obtain

$$f(T) - f(0) = \phi. \quad (6)$$

If we assume that the total wave function rotates by 2π radian for this T which results in a cyclic motion of every state vector of the Hilbert space $\mathcal{H}$, then we will have another condition

$$f(T) - f(0) = 2\pi. \quad (7)$$

Mere satisfaction of condition (7) will ensure the single valuedness of the wave function in the usual Hilbert space $\mathcal{H}$. Again from (4 and 5) we have

$$H|\psi(t)\rangle = i\hbar \left(\frac{d}{dt} \exp(if(t))|\tilde{\psi}(t)\rangle \right) \quad (8)$$

or,

$$H|\psi(t)\rangle = -\hbar \frac{df}{dt} \exp(if(t))|\tilde{\psi}(t)\rangle + i\hbar \exp(if(t)) \frac{d}{dt} |\tilde{\psi}(t)\rangle \quad (9)$$

Since $\psi(t)$ is normalized so we have

$$-\frac{df}{dt} = \frac{1}{\hbar} \langle \psi(t) | H | \psi(t) \rangle - \langle \tilde{\psi}(t) | i \frac{d}{dt} | \tilde{\psi}(t) \rangle. \quad (10)$$

Now integrating equation (10) from 0 to T and using (6) we have

$$\phi = -\frac{1}{\hbar} \int_0^T \langle \psi(t) | H | \psi(t) \rangle dt + \int_0^T \langle \tilde{\psi}(t) | i \frac{d}{dt} | \tilde{\psi}(t) \rangle dt = \gamma + \beta \quad (11)$$

where

$$\gamma = -\frac{1}{\hbar} \int_0^T \langle \psi(t) | H | \psi(t) \rangle dt \quad (12)$$

is proportional to the time integral of the expectation value of the Hamiltonian. It is dynamical in origin and hence called dynamical phase. Now if we subtract the dynamical phase part γ from the total phase ϕ then we will get the geometric phase or the Aharonov Anandan phase

$$\beta = \int_0^T \langle \tilde{\psi}(t) | i \frac{d}{dt} | \tilde{\psi}(t) \rangle dt. \quad (13)$$

It is clear that the same $|\tilde{\psi}(t)\rangle$ can be chosen for infinitely many possible curves C by appropriate choice of $f(t)$. This phase is geometrical in the sense that it does not depend on the Hamiltonian responsible for the evolution. Moreover it is independent of the parameterization and redefinition of phase of $|\psi(t)\rangle$. Hence, β depends only on the evolution of the shadow of $|\psi(t)\rangle$ in projective Hilbert space P and is a pure geometrical entity [3-7].

3 Quasi exactly solvable anharmonic oscillator:

In the introduction we have stated that the QES anharmonic Bose oscillators are very common in Nature. Therefore, a generalized treatment for the geometric phase of the QES anharmonic Bose oscillator may find its application in different directions of physics. Keeping these facts in mind, we will study the possibilities of observing Aharonov-Anandan phase for an QES anharmonic Bose oscillator potential in general.

The Hamiltonian of a generalized QES anharmonic Bose oscillator defined by Dolya and Zaslavaskii [6] is

$$\begin{aligned} H &= H_0 + V \\ &= \sum_{p=1}^{p_0} \epsilon_p (a^\dagger a)^p + \sum_{s=0}^{s_0} A_s \left[(a^\dagger a)^s a^2 + a^{\dagger^2} (a^\dagger a)^s \right] \end{aligned} \quad (14)$$

where a and $a^\dagger$ are the usual annihilation and creation operators respectively. Here we have chosen to work in units in which $m = 1$, $\omega = 1$, $\hbar = 1$. Details of the quasi exactly solvability criterion will be available in the work of Dolya and Zaslavaskii [6]. Now the total eigen state of the unperturbed Hamiltonian H_0 can be written as

$$|\psi(0)\rangle = \sum_n C_n |n\rangle \quad (15)$$

where the coefficients C_n 's depend on the initial conditions. The total wave function of the Hamiltonian H satisfying the Schrödinger equation (4) is

$$\begin{aligned} |\psi(t)\rangle &= \sum_n C_n \exp(-iHt) |n\rangle \\ &= \sum_n C_n \exp \left(\sum_{p=1}^{p_0} -i\epsilon_p (a^\dagger a)^p t + \sum_{s=0}^{s_0} -iA_s \left[(a^\dagger a)^s a^2 + a^{\dagger^2} (a^\dagger a)^s \right] t \right) |n\rangle. \end{aligned} \quad (16)$$

For the calculation of the geometric phase, we have to chose the single-valued state $|\tilde{\psi}(t)\rangle$. This can be easily done by choosing a $f(t)$ which simultaneously satisfies equation (3 and 7). The simplest choice is $f(t) = \lambda t$, where $\lambda = \sum_{s=0}^{s_0} A_s$. Now with that choice we have

$$\begin{aligned} |\tilde{\psi}(t)\rangle &= \exp(-if(t))|\psi(t)\rangle \\ &= \sum_n C_n \exp\left(\sum_{p=1}^{p_0} -i\epsilon_p (a^\dagger a)^p t + \sum_{s=0}^{s_0} -iA_s \left(1 + \left[(a^\dagger a)^s a^2 + a^{\dagger 2} (a^\dagger a)^s\right]\right) t\right) |n\rangle \end{aligned} \quad (17)$$

and

$$f(T) = \lambda T = 2\pi \quad (18)$$

Therefore, from equations (13,17 and 18) a compact expression for the Aharonov Anandan phase of a QES Bose anharmonic system can be obtained as

$$\begin{aligned} \beta &= \int_0^T \langle \tilde{\psi}(t) | i \frac{d}{dt} | \tilde{\psi}(t) \rangle dt \\ &= \int_0^T \sum_k \sum_n \langle k | C_k^* C_n \exp\left(\sum_{p=1}^{p_0} i\epsilon_p (a^\dagger a)^p t + \sum_{s=0}^{s_0} iA_s \left(1 + \left[(a^\dagger a)^s a^2 + a^{\dagger 2} (a^\dagger a)^s\right]\right) t\right) \\ &\times \left(\sum_{p=1}^{p_0} \epsilon_p (a^\dagger a)^p + \sum_{s=0}^{s_0} A_s \left(1 + \left[(a^\dagger a)^s a^2 + a^{\dagger 2} (a^\dagger a)^s\right]\right) t\right) \\ &\times \exp\left(\sum_{p=1}^{p_0} -i\epsilon_p (a^\dagger a)^p t + \sum_{s=0}^{s_0} -iA_s \left(1 + \left[(a^\dagger a)^s a^2 + a^{\dagger 2} (a^\dagger a)^s\right]\right) t\right) |n\rangle dt \\ &= \int_0^T \sum_k \sum_n \langle k | C_k^* C_n \left(\sum_{p=1}^{p_0} \epsilon_p (a^\dagger a)^p + \sum_{s=0}^{s_0} A_s \left(1 + \left[(a^\dagger a)^s a^2 + a^{\dagger 2} (a^\dagger a)^s\right]\right)\right) |n\rangle dt \\ &= 2\pi + T \sum_n |C_n|^2 \sum_{p=1}^{p_0} \epsilon_p (n)^p + T \sum_n C_{n-2}^* C_n \sum_{s=0}^{s_0} A_s (n-2)^s \sqrt{n(n-1)} \\ &+ T \sum_n C_{n+2}^* C_n \sum_{s=0}^{s_0} A_s (n)^s \sqrt{(n+1)(n+2)}. \end{aligned} \quad (19)$$

The quantum nature of the geometric phase is manifested here through the discrete sum appearing in the expression (19) and the state dependence of the phase appears through the coefficients $C_k^* C_n$. Now let us think of some special situations of physical interest.

3.1 Input is an intense laser beam.

Let us now consider the case where an intense electromagnetic field having Poissonian statistics ($C_n = \exp(-\frac{|\alpha|^2}{2}) \frac{\alpha^n}{\sqrt{n!}}$) interacts with a system characterized by the Hamiltonian (14). With the help of equations (19) the geometric phase β for this physical system can be written as

$$\begin{aligned} \beta &= 2\pi + T \sum_n |C_n|^2 \sum_{p=1}^{p_0} \epsilon_p (n)^p + T \sum_n C_{n-2}^* C_n \sum_{s=0}^{s_0} A_s (n-2)^s \sqrt{n(n-1)} \\ &+ T \sum_n C_{n+2}^* C_n \sum_{s=0}^{s_0} A_s (n)^s \sqrt{(n+1)(n+2)} \\ &= 2\pi + T \exp(-|\alpha|^2) \sum_n \frac{|\alpha|^{2n}}{n!} \left(\sum_{p=1}^{p_0} \epsilon_p (n)^p + \sum_{s=0}^{s_0} 2|\alpha|^2 A_s (n)^s \cos(2\theta)\right). \end{aligned}$$

Photon Statistics	$ C_n ^2$
Sub-Poissonian	$\frac{N!}{(N-n)!n!} p^n (1-p)^{N-n} \quad (0 \leq p \leq 1)$
Poissonian	$\exp(- \alpha ^2) \frac{ \alpha ^{2n}}{n!}$
Super-Poissonian	$\frac{(n+W-1)!}{(W-1)!n!} q^n (1-q)^W \quad (0 \leq q \leq 1)$

table1

where $\alpha = |\alpha| \exp(i\theta)$. Here we observe that the geometric phase depends on the phase of the initial laser beam. The presence of the off-diagonal terms in the interaction Hamiltonian is manifested through the presence of θ dependent terms in the expression for the geometric phase β .

4 Discussions and Conclusions:

In the present work we have seen that the geometric phase appears in QES Bose systems and depends on the photon statistics of the input field through C_n 's (see table 1). We have also observed that the presence of the off-diagonal terms in the interaction Hamiltonian is manifested through the presence of θ dependent terms in the expression for the geometric phase β . From these facts we can conclude two things, firstly, one should always have to take care of the possibilities of appearance of geometric phase while working with QES Bose systems and, secondly, geometric phase can be tuned because it depends strongly on the phase of the input field which can be tuned.

Acknowledgement: Author is thankful to CSIR, India for a senior research fellowship.

References:

1. Perinova V, Luks A and Perina J 1998 *Phases in Optics* (New Jersey: World Scientific)
2. Berry M V 1984 *Proc. R. Soc A* **392** 45.
3. Aharonov Y and Anandan J 1987 *Phys. Rev. Lett.*, **58** 1593.
4. Joshi A, Pati A and Banerjee A 1994 *Phys. Rev. A* 49 5131.
5. Blais A and Tremblay A M S *Preprint* quant-ph/0105006.
6. Dolya S N and Zaslavskii O B 2000 *J. Phys. A* **33** L369.
7. Moore D J 1991 *Phys. Rep.* **210** 1.
8. Dass T and Sharma S K 1998 *Mathematical Methods in Classical and Quantum Physics* (Hyderabad: University Press)